PNU-NTG-06/2004

Magnetic susceptibility of the QCD vacuum

Hyun-Chul Kim,[1,*] M. Musakhanov,[1,2,†] and M. Siddikov[2,3,‡]

[1] *Department of Physics and Nuclear Physics & Radiation Technology Institute (NuRI),*
Pusan National University, 609-735 Busan, Republic of Korea
[2] *Theoretical Physics Department, National University*
of Uzbekistan, Tashkent 700174, Uzbekistan
[3] *Institut für Theoretische Physik II,*
Ruhr-Universität Bochum, D–44780 Bochum, Germany

(Dated: November 2004)

arXiv:hep-ph/0411181v1 12 Nov 2004

Abstract

We investigate the magnetic susceptibility of the QCD vacuum, based on the instanton vacuum. Starting from the instanton liquid model for the instanton vacuum, we derive the light-quark partition function $Z[V, T, \hat{m}]$ in the presence of the current quark mass $\hat{m}$ as well as the external Abelian vector and tensor fields. We calculate a two-point correlation function relevant for the magnetic susceptibility and derive it beyond the chiral limit. We obtain for the different flavors the following magnetic susceptibility: $\chi_{u,d}\langle i\psi^{\dagger}_{u,d}\psi_{u,d}\rangle_0 \sim 40 \sim 45\,\mathrm{MeV}$, while $\chi_s\langle i\psi^{\dagger}_s\psi_s\rangle_0 \simeq 6 \sim 10\,\mathrm{MeV}$ with the quark condensate $\langle i\psi^{\dagger}\psi\rangle_0$.

PACS numbers:
Keywords:

---

[*] Electronic address: hchkim@pusan.ac.kr
[†] Electronic address: musakhanov@pusan.ac.kr
[‡] Electronic address: Marat.Siddikov@tp2.ruhr-uni-bochum.de

1. The QCD vacuum is one of the most complicated states filled not only by perturbative but also strong nonperturbative fluctuations. In particular, the condensates of the quarks and gluons characterize its nonperturbative aspect. The quark condensate plays a role of an order parameter associated with spontaneous chiral symmetry breaking (SχSB) which is probably one of the most important quantities for low-energy hadronic phenomena.

The presence of a constant external electromagnetic field $F_{\mu\nu}$ induces another type of condensates leading to the nonzero magnetic susceptibility χ_f of the QCD vacuum defined as:

$$\langle 0|\psi_f^\dagger \sigma_{\mu\nu}\psi_f|0\rangle_F = e_f\,\chi_f\,\langle i\psi_f^\dagger\psi_f\rangle_0\,F_{\mu\nu}, \tag{1}$$

where e_f denotes the quark electric charge, f is the corresponding flavor. It is natural to have the quark condensate $\langle i\psi_f^\dagger\psi_f\rangle_0$ in the chiral limit as a normalization factor in the right-hand side of Eq. (1), since SχSB is responsible for these quantities, i.e. the quark condensate and magnetic susceptibility. In a recent paper [1] it was suggested that the magnetic susceptibility χ_f may be measured in the exclusive photoproduction of hard dijets $\gamma + N \to (\bar{q}q) + N$. The value of χ_f was predicted by using the method of the QCD sum rule and vector dominance [2, 3, 4]: $\chi_f\langle i\psi_f^\dagger\psi_f\rangle_0 = 40 \sim 70\,\mathrm{MeV}$ at the scale of $1\,\mathrm{GeV}$.

In the present work, we want to investigate the magnetic susceptibility χ_f of the QCD vacuum within the framework of the instanton vacuum. Since the instanton vacuum explains SχSB naturally via quark zero modes, it may provide a good framework to study the χ_f. Moreover, there are only two parameters in this approach, namely the average instanton size $\rho \approx \frac{1}{3}\,\mathrm{fm}$ and average inter-instanton distance $R \approx 1\,\mathrm{fm}$. The normalization scale of this approach can be defined by the average size of instantons and is approximately equal to $\rho^{-1} \approx 0.6\,\mathrm{GeV}$. The values of the ρ and R were estimated many years ago phenomenologically by Ref. [5] as well as theoretically by Ref. [6]. Furthermore, it was recently confirmed in various lattice simulations of the QCD vacuum [7, 8, 9]. Very recent lattice calculations of the quark propagator [10, 11] are in a remarkable agreement with that of Ref. [6].

The magnetic susceptibility of the QCD vacuum was estimated already in Ref. [20], however, done in the chiral limit and without the consideration of the current conservation. Since we want to investigate it beyond the chiral limit, we first need to extend Refs. [12, 13, 14] and Lee-Bardeen formula [15] for a low-frequency ($p \ll \rho^{-1}$) part of the quark determinant:

$$\mathrm{Det}_{\mathrm{low}} = \det B,\;\; B_{ij} = i\hat{m}\delta_{ij} + a_{ji},\;\; a_{-+} = -\langle \Phi_{-,0}|i\partial\!\!\!/|\Phi_{+,0}\rangle, \tag{2}$$

taking into account the current quark mass. Here, $\hat{m}$ denotes a current quark mass, which is assumed to be small, a_{ij} stands for the overlapping matrix element of the quark zero modes $\Phi_{\pm,0}$ generated by instantons (antiinstantons). In general, we define $\Phi_{\pm,n}$ by the following equation:

$$(i\partial\!\!\!/ + A\!\!\!/_{\pm})\Phi_{\pm,n} = \lambda_n \Phi_{\pm,n}, \tag{3}$$

where $A_{\mu,\pm}$ is an instanton (antiinstanton) field and $\lambda_0 = 0$ by convention. The matrix element a_{ij} is nonzero only between instantons and antiinstantons and vice versa due to specific chiral properties of the zero modes.

The overlapping of the quark zero modes allows quarks to propagate through the instanton-antiinstanton medium by jumping from one instanton to another one. Hence, $\mathrm{Det}_{\mathrm{low}}$ is reduced to the determinant of the finite matrix B in the subspace of *zero modes.*

This result was reproduced by Refs. [13, 16] in different ways. It is well known that the fermionic determinant is proportional to $m^{|N_+-N_-|}$ and strongly suppresses the fluctuations of $|N_+ - N_-|$. Therefore, we can take $N_+ = N_- = N/2$ in the final formulas.

In order to calculate the magnetic susceptibility, we have to find the light-quark partition function $Z[V, T, \hat{m}]$ in the presence of the Abelian external vector V_μ and antisymmetric tensor $T_{\mu\nu}$ fields. So, we first calculate a zero-mode approximated quark propagator $\tilde{S}$ in the instanton ensemble and in the presence of these external fields. Using this propagator, we are able to compute a low-frequency part of the quark determinant $\widetilde{\mathrm{Det}}_{\mathrm{low}}$ and arrive at the extension of Eq. (2) for the non-zero $\hat{m}$ with the external Abelian vector V and tensor T fields.

The smallness of the packing parameter $\pi(\frac{\rho}{R})^4 \approx 0.1$ makes it possible to average the determinant over collective coordinates of instantons with fermionic quasiparticles, i.e. constituent quarks ψ introduced. The averaged determinant turns out to be the light-quark partition function $Z[V, T, \hat{m}]$ which is a functional of V and T and can be represented by a functional integral over the constituent quark fields with the gauged effective chiral action $S[\psi^\dagger, \psi, V, T]$. However, it is not trivial to make the action gauge-invariant due to the nonlocality of the quark-quark interactions generated by instantons. In the previous paper [17], it was demonstrated how to gauge the nonlocal effective chiral action in the presence of the external electromagnetic field. The gauged action was successfully tested by showing that the low-energy theorem of the axial anomaly related to the process $G\tilde{G} \to \gamma\gamma$ (see also [18, 19]) is satisfied.

In the present work we refine the derivation of the gauged effective chiral action and apply it to the calculations of the magnetic susceptibility χ_f of the QCD vacuum, taking into account both current quark mass and nonlocal currents.

2. It is clear that the magnetic susceptibility χ_f is related to the two-point correlation function of the quark tensor and vector currents at the zero momentum. In order to calculate this correlation function, we need to treat the light-quark partition function $Z[V, T, \hat{m}]$ as a functional of external abelian vector V_μ and tensor $T_{\mu\nu}$ fields. The relevant total quark determinant

$$\widetilde{\mathrm{Det}} := \mathrm{Det}(i\partial\!\!\!/ + A\!\!\!/ + eV\!\!\!/ + \sigma_{\mu\nu}T_{\mu\nu} + i\hat{m}) \tag{4}$$

can be splitted into two parts which correspond to low and high momenta, respectivley, with respect to some auxiliary parameter M_1 lying inside an interval, $R^{-1} \ll M_1 \ll \rho^{-1}$: $\widetilde{\mathrm{Det}} = \widetilde{\mathrm{Det}}_{\mathrm{low}} \times \widetilde{\mathrm{Det}}_{\mathrm{high}}$ [12]. $\widetilde{\mathrm{Det}}_{\mathrm{high}}$ comes from fermion modes with Dirac eigenvalues from the interval M_1 to the Pauli–Villars mass M, and $\widetilde{\mathrm{Det}}_{\mathrm{low}}$ accounts for eigenvalues less than M_1. The product of these determinants is independent of the scale M_1. However, we can deal with them only approximately. The high-momentum part $\widetilde{\mathrm{Det}}_{\mathrm{high}}$ can be written as a product of the determinants in the field of individual instantons, while the low-momentum one $\widetilde{\mathrm{Det}}_{\mathrm{low}}$ has to be treated approximately, would-be zero modes being only taken into account. In Ref. [12], it was demonstrated that the $\widetilde{\mathrm{Det}}_{\mathrm{low}}$ depends weakly on M_1 in the wide range, which serves as a check for this approximation. Based on this prescription, we now calculate the low-momentum part of the quark determinant in the presence of instantons as well as external V_μ, $T_{\mu\nu}$ fields, and then average it over the collective coordinates of the instantons in order to find the effective low-energy QCD partition function as a functional of the external field $Z[V, T, \hat{m}]$.

We first define the total quark propagator $\tilde{S}$ in the presence of the instanton ensemble A and external fields V, T and the quark propagator $\tilde{S}_i$ with a single instanton A_i as well as

V and T:

$$\tilde{S} = \frac{1}{i\partial\!\!\!/ + gA\!\!\!/ + eV\!\!\!/ + \sigma\cdot T + i\hat{m}}, \quad \tilde{S}_i = \frac{1}{i\partial\!\!\!/ + gA\!\!\!/_i + eV\!\!\!/ + \sigma\cdot T + i\hat{m}}, \tag{5}$$

where $\sigma\cdot T = \sigma_{\mu\nu}T_{\mu\nu}$. We assume that the total instanton field A may be approximated as a sum of the single instanton fields, $A = \sum_{i=1}^{N} A_i$, which is justified with the above-mentioned average size of instantons $\rho \approx 1/3\,\mathrm{fm}$ and average inter-instanton distance $R \approx 1\,\mathrm{fm}$. Defining the quark propagator $\tilde{S}_0$ with external fields V and T and the free one S_0 as follows:

$$\tilde{S}_0 = \frac{1}{i\partial\!\!\!/ + eV\!\!\!/ + \sigma\cdot T + i\hat{m}}, \quad S_0 = \frac{1}{i\partial\!\!\!/ + i\hat{m}}, \tag{6}$$

we can expand the quark propagator $\tilde{S}$ with respect to a single instanton:

$$\tilde{S} = \tilde{S}_0 + \sum_i (\tilde{S}_i - \tilde{S}_0) + \sum_{i\neq j} (\tilde{S}_i - \tilde{S}_0)\tilde{S}_0^{-1}(\tilde{S}_j - \tilde{S}_0) + \cdots . \tag{7}$$

In order to specify the gauge dependence, we rewrite $\tilde{S}_i$ and S_0 in the following form:

$$\begin{aligned} \tilde{S}_i &= L_i S_i' L_i^{-1}, \quad S_i' = \frac{1}{i\partial\!\!\!/ + gA\!\!\!/_i + eV\!\!\!/' + \sigma\cdot T + i\hat{m}}, \\ \tilde{S}_0 &= L_i S_0' L_i^{-1}, \quad S_{0i}' = \frac{1}{i\partial\!\!\!/ + eV\!\!\!/' + \sigma\cdot T + i\hat{m}}, \end{aligned} \tag{8}$$

where $eV\!\!\!/' = L^{-1}(i\partial\!\!\!/ + eV\!\!\!/)L$ and the gauge connection L can be written as the path-ordered exponent:

$$L_i(x, z_i) = \mathrm{P}\exp\left(ie\int_{z_i}^{x} d\xi_\mu V_\mu(\xi) \right), \tag{9}$$

where z_i denotes an instanton position.

In the case of the small quark mass $\hat{m}$, we can expand the quark propagator S_i [12] as follows:

$$S_i(\hat{m}\to 0) = \frac{1}{i\partial\!\!\!/} + \frac{|\Phi_{0i}\rangle\langle\Phi_{0i}|}{i\hat{m}}. \tag{10}$$

While it gives a proper value for the $\langle\Phi_{0i}|S_i(\hat{m}\to 0)|\Phi_{0i}\rangle = \frac{1}{i\hat{m}}$, the second term has wrong chiral properties:

$$S_i(\hat{m}\to 0)|\Phi_{0i}\rangle = \frac{|\Phi_{0i}\rangle}{i\hat{m}} + \frac{1}{i\partial\!\!\!/}|\Phi_{0i}\rangle. \tag{11}$$

However, we may neglect this term as $\hat{m}\to 0$.

If the $\hat{m}$ is not small, then we propose the following approximation:

$$S_i = S_0 + S_0 i\partial\!\!\!/ \frac{|\Phi_{0i}\rangle\langle\Phi_{0i}|}{c_i} i\partial\!\!\!/ S_0, \tag{12}$$

where

$$c_i = -\langle\Phi_{0i}|i\not\partial S_0 i\not\partial|\Phi_{0i}\rangle = i\hat{m}\langle\Phi_{0i}|S_0 i\not\partial|\Phi_{0i}\rangle. \tag{13}$$

The merit of the approximation proposed in Eq.(12) lies in projecting S_i onto the zero-mode:

$$S_i|\Phi_{0i}\rangle = \frac{1}{i\hat{m}}|\Phi_{0i}\rangle,\ \ \langle\Phi_{0i}|S_i = \langle\Phi_{0i}|\frac{1}{i\hat{m}}. \tag{14}$$

Including the external fields V and T, we get

$$\tilde{S}_i = \tilde{S}_{0i} + \tilde{S}_{0i}L_i i\not\partial\frac{|\Phi_{0i}\rangle\langle\Phi_{0i}|}{c_i - b_i}i\not\partial L_i^{-1}\tilde{S}_{0i}, \tag{15}$$

where

$$b_i - c_i = \langle\Phi_{0i}|i\not\partial(L_i^{-1}\tilde{S}_0 L_i)i\not\partial|\Phi_{0i}\rangle. \tag{16}$$

Thus, we can express Eq.(7) as

$$\tilde{S} = \tilde{S}_0 + \tilde{S}_0\sum_{i,j}L_i i\not\partial|\Phi_{i0}\rangle\left(\frac{1}{-D-T}\right)_{ij}\langle\Phi_{0j}|i\not\partial L_j^{-1}\tilde{S}_0, \tag{17}$$

where

$$D_{ij} + T_{ij} = \langle\Phi_{0i}|i\not\partial(L_i^{-1}\tilde{S}_0 L_j)i\not\partial|\Phi_{0j}\rangle,\ \ D_{ij} = (b_i - c_i)\delta_{ij}. \tag{18}$$

Introducing the following function:

$$|\phi_{0i}\rangle = \frac{1}{i\not\partial}L_i i\not\partial|\Phi_{0i}\rangle, \tag{19}$$

which has the same chiral properties as the zero-mode eigenfunction $|\Phi_{0i}\rangle$, and taking the trace over the subspace of the zero-modes, we obtain the following expression:

$$\tilde{\mathrm{Tr}}(\tilde{S} - \tilde{S}_0) = -\sum_{i,j}\langle\phi_{0,j}|i\not\partial\,(\tilde{S}_0^2)\,i\not\partial|\phi_{0,i}\rangle\langle\phi_{0,i}|(\frac{1}{i\not\partial\tilde{S}_0 i\not\partial})|\phi_{0,j}\rangle, \tag{20}$$

where $\tilde{\mathrm{Tr}}$ denotes the trace over the zero-mode subspace only. Bringing now in the matrix

$$\tilde{B}(m)_{ij} = \langle\phi_{0,i}|(i\not\partial\tilde{S}_0 i\not\partial)|\phi_{0,j}\rangle, \tag{21}$$

we get

$$\tilde{\mathrm{Tr}}\int_{M_1}^{m} idm'(\tilde{S}(m') - \tilde{S}_0(m')) = \tilde{\mathrm{Tr}}\int_{\tilde{B}(M_1)}^{\tilde{B}(m)} d\tilde{B}(m')\frac{1}{\tilde{B}(m')} = \tilde{\mathrm{Tr}}\ln\frac{\tilde{B}(m)}{\tilde{B}(M_1)}, \tag{22}$$

from which we find that $\tilde{B} = \tilde{B}(m)$ is equivalent to a mere extension of the Lee-Bardeen matrix B [15] in the presence of the external Abelian fields V, T and current quark mass $\hat{m}$. Thus, we derive for arbitrary flavor N_f

$$\tilde{B}^f_{ij} = \langle\Phi_{0i}|i\not\partial L^\dagger_{fi}\,\tilde{S}_{0f}L_{fj}i\not\partial|\Phi_{0j}\rangle \tag{23}$$

$$\tilde{S}_{0f} = \frac{1}{i\partial\!\!\!/ + e_f V\!\!\!/ + \sigma\cdot T + i m_f},\ \ L_{fi} = \mathrm{P}\exp\left(ie_f\int_{z_i}^{x} d\xi_\mu V_\mu(\xi)\right). \tag{24}$$

If we introduce Grassmanian variables $\bar{\Omega}_i^f$ and Ω_j^g, then we can express the fermionic determinant of B as a functional integral over fermion fields:

$$\begin{aligned}\det\tilde{B} \ &= \ \int d\Omega d\bar{\Omega}\exp\left[(\bar{\Omega}\tilde{B}\Omega),\ (\bar{\Omega}\tilde{B}\Omega)\right] = \bar{\Omega}_i^f\langle\phi_{0,i}|i\partial\!\!\!/\tilde{S}_{0,f}i\partial\!\!\!/|\phi_{0,j}\rangle\Omega_j^f \\ &= \ (\mathrm{Det}(i\partial\!\!\!/ + eV\!\!\!/ + \sigma\cdot T + i\hat{m}))^{-1}\int\prod_{i,f} d\Omega_i^f d\bar{\Omega}_i^f D\psi_f D\psi_f^\dagger \\ &\times \ \mathrm{e}^{\int dx\left\{\psi_f^\dagger(x)(i\partial\!\!\!/ + e_f V\!\!\!/ + \sigma\cdot T + i m_f)\psi_f(x) + \bar{\eta}_i^f(x)\psi_f(x) + \psi_f^\dagger(x)\eta_i^f(x)\right\}},\end{aligned} \tag{25}$$

where the source fields η_i^f and $\bar{\eta}_j^f$ are defined as:

$$\bar{\eta}_i^f = \bar{\Omega}_i^f\langle\phi_{0,i}|i\partial\!\!\!/,\ \ \eta_j^f = i\partial\!\!\!/|\phi_{0,j}\rangle\Omega_j^f. \tag{26}$$

The integration over the Grassmanian variables Ω and $\bar{\Omega}$ provides finally the low-frequency part of the quark determinant in the following form:

$$\tilde{\mathrm{Det}}_{\mathrm{low}} = \det\tilde{B} = (\det(i\partial\!\!\!/ + eV\!\!\!/ + \sigma\cdot T + i\hat{m}))^{-1}\int\prod_f D\psi_f D\psi_f^\dagger \tag{27}$$

$$\times\ \mathrm{e}^{\left(\int d^4x\psi_f^\dagger(i\partial\!\!\!/ + e_f V\!\!\!/ + \sigma\cdot T + i m_f)\psi_f\right)}\prod_f\left\{\prod_{+}^{N_+} V_{+,f}[\psi_f^\dagger,\psi_f]\prod_{-}^{N_-} V_{-,f}[\psi_f^\dagger,\psi_f]\right\},$$

where

$$\tilde{V}_{\pm,f}[\psi_f^\dagger,\psi_f] = \int d^4x\left(\psi_f^\dagger(x)\,L_f(x,z)i\partial\!\!\!/\Phi_{\pm,0}(x;\xi_\pm)\right)\int d^4y\left(\Phi_{\pm,0}^\dagger(y;\xi_\pm)(i\partial\!\!\!/ L_f^+(y,z)\psi_f(y)\right) \tag{28}$$

It is obvious that $\tilde{V}_{\pm,f}[\psi^\dagger,\psi]$ indicates the nonlocal interaction between constituent quarks generated by instantons. Since the range of the integration in Eq.(28) is truncated at ρ, which is defined by zero-mode functions $\Phi_{\pm,0}$, the range of the nonlocality is determined by ρ.

Note that the external vector field V_μ gauges not only the kinetic term of the effective action but also its interaction one $\tilde{V}_{\pm,f}[\psi_f^\dagger,\psi_f]$ in Eq. (28). The reason is clear: Since the instanton-induced interaction is nonlocal, the gauge connection L_f must be attached to each fermionic line so that the effective action may be gauge-invariant. Unfortunately, we encounter an ambiguity arising from this gauge connection, since it is path-dependent in general. However, we will show now that the most preferable path can be found in the present case.

3. In order to solve the problem of gauge dependence due to the nonlocal interaction induced by instantons, we have employed the gauge connection L_f. Though it causes in general some arbitrariness due to its path dependence, we can prove that such dependence can be minimized in the present work. To be more specific, we consider the extended zero mode:

$$(iD\!\!\!\!/ + V\!\!\!/)|\tilde{\Phi}_0\rangle = 0,\ \ iD\!\!\!\!/ = i\partial\!\!\!/ + A\!\!\!/,\ \ |\tilde{\Phi}_0^{(1)}\rangle = |\Phi_0\rangle - S_{NZ}V\!\!\!/|\Phi_0\rangle, \tag{29}$$

where A denotes an instanton field located at z, $|\tilde{\Phi}_0^{(1)}\rangle$ stands for the solution to order $\mathcal{O}(V)$, and S_{NZ} the well-known non-zero mode of the propagator in the instanton field (see the review [5] and references therein). Putting the gauge connection $L(x,z)$ defined in Eq. (9) into Eq. (29), we obtain:

$$(i\not{D} + \not{V}')|\Phi_0'\rangle = 0, \;\; V_\mu' = L^{-1}(i\partial_\mu + V_\mu)L, \;\; |\Phi_0'\rangle = L^{-1}(x,z)|\tilde{\Phi}_0\rangle. \tag{30}$$

$i\not{D}\,|\Phi_0\rangle = 0$ and $S_{NZ}\, i\not{D} = 1 - |\Phi_0\rangle\langle\Phi_0|$ being used, the solution $|\Phi_0^{(1)'}\rangle$ to order $\mathcal{O}(V)$ in Eq. (30) can be reduced to the corresponding solution $|\tilde{\Phi}_0^{(1)}\rangle$ to order $\mathcal{O}(V)$ without any trace of the path dependence arising from the gauge connection L.

However, if the zero-mode approximation $S_{NZ} \approx \frac{1}{i\not{\partial}} = S_{00}$ is used to find the solution to order $\mathcal{O}(V)$

$$|\Phi_{00}^{(1)'}\rangle = |\Phi_0\rangle - S_{00}\not{V}'|\Phi_0\rangle, \tag{31}$$

then applying the inverse gauge connection L^{-1} to Eq.(31) leads to

$$|\tilde{\Phi}_{00}^{(1)}\rangle = |\Phi_0\rangle - S_{00}\not{V}|\Phi_0\rangle + S_{00}(i\int d\xi_\mu V_\mu(\xi))|\Phi_0\rangle, \tag{32}$$

which depends on the path we choose explicitly. Thus, it is the best way that we choose the path minimizing $\|\tilde{\Phi}_0^{(1)} - \tilde{\Phi}_{00}^{(1)}\|^2$. Expressing the external vector field in a Fourier form:

$$V_\mu(x) = \int d^4q V_\mu(q) e^{iq\cdot x}, \tag{33}$$

we can parameterize the path from x to z as follows:

$$\xi_\mu(x,z,t) = z_\mu + (x-z)_\mu t + p_\mu(x,z,t) \tag{34}$$

with parameter $t \in (0,1)$ and boundary conditions $p_\mu(t=0) = p_\mu(t=1) = 0$. It is the most optimized path we can take, since p_μ should depend only on the path difference $x-z$ in order to satisfy the conservation of the linear momenta. More explicitly, we write down the path in the following form:

$$\xi_\mu(x,z,t) = z_\mu + (x-z)_\mu t + p_\mu t(1-t). \tag{35}$$

Then we have

$$i\int d\xi_\mu V_\mu(\xi) = i\int d^4q V_\mu(q) e^{iq\cdot z}\int_0^1 dt[(x-z)_\mu + (1-2t)p_\mu] e^{iq\cdot[(x-z)t + ipt(1-t)]}. \tag{36}$$

In principle, we can express Eq. (36) in terms of the error function. However, we shall consider a simple case of $q\rho \ll 1$ so that we can expand the vector current with respect to $(q\cdot p)$ and $q\cdot(x-z)$:

$$V_\mu(x) = \int d^4q V_\mu(q) e^{iq\cdot x} \approx \int d^4q V_\mu(q) e^{iq\cdot z}\left[1 + iq\cdot(x-z) - \frac{(q\cdot(x-z))^2}{2}\right] \tag{37}$$

Substituting this expansion into Eqs. (29) and (32) and carrying out some tedious but straightforward calculations, we obtain

$$\|\tilde{\Phi}_0^{(1)} - \tilde{\Phi}_{00}^{(1)}\|^2 \approx C\rho^2 N_f \int d^4q V^2(q)(q\cdot p)^2 + \cdots \tag{38}$$

where the constant $C = +0.0063 > 0$ and higher-order terms are neglected. Therefore, we can conclude that the minimum is achieved by taking the straight-line path that will be employed in the sequel.

Moreover, since we are only interested in the case of a constant electromagnetic field strength $F_{\mu\nu}$, we can even show that the magnetic susceptibility does not depend at all on the path. Let us consider the following difference:

$$\delta\left(\int_z^x d\xi_\mu V_\mu(\xi)\right) := \int_C d\xi_\mu V_\mu(\xi) - \int_{s-l} d\xi_\mu V_\mu(\xi), \tag{39}$$

where $\int_C d\xi_\mu V_\mu(\xi)$ is the line integral along the path C parameterized by $\xi_\mu(x,z,t) = z_\mu + (x-z)_\mu t + a_\mu(x-z,t)$ and $\int_{s-l} d\xi_\mu V_\mu(\xi)$ is calculated along the straight-line. This difference is reducible to the integral over the surface between these two contours due to Stoke's theorem:

$$\delta\left(\int_z^x d\xi_\mu V_\mu(\xi)\right) := \oint d\xi_\mu V_\mu(\xi) = \oint dS_{\mu\nu} F_{\mu\nu}(\xi) = F_{\mu\nu}\int d\xi_\mu a_\nu(\xi), \tag{40}$$

where the integral $\int d\xi_\mu a_\nu(\xi)$ depends only on a distance $(x-z)$ but not on x and z separately due to translational invariance. Denoting $G_{\mu\nu}(x-z) \equiv \int d\xi_\mu a_\nu(\xi) F(x-z)$, where $F(x-z)$ represents the form factor F in coordinate space, we find that the change of the magnetic susceptibility $\delta\chi_f$ due to the deformation of the contour turns out to vanish:

$$F_{\lambda\rho}\mathrm{Tr}[S(p)G_{\rho\lambda}(p)F(p)S(p)\sigma_{\mu\nu} + S(p)F(p)G_{\rho\lambda}(p)S(p)\sigma_{\mu\nu}] = 0. \tag{41}$$

Thus, we are now allowed to take any path without changing the final results of the magnetic susceptibility.

4. We are now in a position to derive the relevant partition function for the magnetic susceptibility of the QCD vacuum. We first have to average the low-frequency part of the quark determinant $\tilde{\det}_{\rm low}$ over collective coordinates $\xi_\pm$. Having exponentiated and bosonized the interaction terms [12, 13, 17, 25, 26], we arrive at the effective chiral action with external fields pertinent to the magnetic susceptibility. Since the density of the instanton medium ($\pi^2\left(\frac{\rho}{R}\right)^4 \approx 0.1$) is low, we are able to average over positions and orientations of the instantons independently. In the next step we have to introduce auxiliary fields, i.e. the couplings $\lambda_\pm$ for exponentiation and meson fields $\Phi_\pm(x)$ for bosonization, respectively. Having integrated over fermion fields, we end up with the partition function in the presence of the external vector and tensor fields:

$$Z[V,T,\hat{m}] = \int d\lambda_+ d\lambda_- D\Phi_+ D\Phi_- \exp\left(-S[\lambda_\pm,\Phi_\pm]\right), \tag{42}$$

where

$$\begin{aligned} S[\lambda_\pm,\Phi_\pm] &= -\sum_\pm\left[N_\pm\ln\left\{\left(\frac{4\pi^2\rho^2}{N_c}\right)^{N_f}\frac{N_\pm}{V\lambda_\pm}\right\}-N_\pm\right]+S_\Phi+S_\psi, \qquad (43)\\ S_\Phi &= \int d^4x\sum_\pm(N_f-1)\lambda_\pm^{-\frac{1}{N_f-1}}(\det\Phi_\pm(x))^{\frac{1}{N_f-1}},\\ S_\psi &= -\mathrm{Tr}\ln((\not{P}+\sigma\cdot T+i\hat{m}+iF(P)\sum_\pm\Phi_\pm(x)\frac{1\pm\gamma_5}{2})F(P)(\not{P}+\sigma\cdot T+i\hat{m})^{-1}). \end{aligned}$$

Here, $P_\mu = p_\mu + eV_\mu$, $p_\mu = i\partial_\mu$ and Tr stands for the functional trace, i.e. $\int d^4x\mathrm{tr}_c\mathrm{tr}_D\mathrm{tr}_f$ and $F(k)$ denotes the quark form factor generated by the fermionic zero modes and has an explicit form as follows [12]:

$$F(k\bar{\rho}) = 2z\left(I_0(z)K_1(z)-I_1(z)K_0(z)-\frac{1}{z}I_1(z)K_1(z)\right), \qquad (44)$$

where I_0, I_1, K_0, and K_1 are the modified Bessel functions, z is defined as $z = k\bar{\rho}/2$. When k goes to infinity, this zero-mode form factor $F(k\bar{\rho})$ has the following asymptotic behavior:

$$F(k\bar{\rho}) \longrightarrow \frac{6}{(k\bar{\rho})^3}. \qquad (45)$$

In fact, the nonlocal effective quark-meson interaction can be understood without relying on the instanton vacuum [23]. In those cases, the momentum-dependent quark mass can be interpreted as a nonlocal regularization in Euclidean space. Hence, various types of the $F(k)$ as a regulator with the regularization parameter $\Lambda \sim 1/\bar{\rho}$ has been used by different authors. For example, the dipole-type $F(k)$ is adopted in the study of the pion wave function [20], while the Gaussian is employed in Ref. [24]. Hence, we utilize in this work the dipole and exponential types of the form factors in addition to the instanton-induced zero-mode form factor:

$$F(k) = \left\{\begin{array}{ll}\Lambda^2/(\Lambda^2+k^2) & \text{(dipole)},\\ \exp(-k^2/\Lambda^2) & \text{(Gaussian)}.\end{array}\right. \qquad (46)$$

The saddle-point approximation of the integrals in the partition function given in Eq. (42) leads to the equation for the effective quark mass, which is in general expressed by the functional of the external fields V and T. However, since we finally need the term of order $\mathcal{O}(VT)$, we can consider the saddle-point approximation without external fields:

$$\Phi_{\pm,fg} = \Phi_{\pm,fg}(0) = M_f\delta_{fg}, \quad \lambda_\pm = \lambda = \frac{2V}{N}\prod_f M_f, \qquad (47)$$

which satisfies the following saddle-point equation

$$\frac{N}{V} = 4N_c\int\frac{d^4k}{(2\pi)^4}\frac{M_fF^2(k)(m_f+M_fF^2(k))}{k^2+(m_f+M_fF^2(k))^2}. \qquad (48)$$

Equation (48) contains the momentum-dependent dynamic quark mass $M_f(k) = M_fF^2(k)$. Note that the dynamic quark mass M_f is a decreasing function of the current quark mass

m_f [25, 26] with meson loops neglected. At small m_f, we may write $M_f = M_0 + \gamma m_f$ with $M_0 \approx 360$ MeV and $\gamma \approx -2$[1].

Having derived the partition function in Eq. (43), we are now ready for the calculation of the magnetic susceptibility χ_f from the instanton vacuum within our framework. In order to evaluate χ_f, we have to compute the following correlation function via the functional derivative with respect to the external tensor field:

$$\frac{\delta}{\delta T_{\mu\nu}} \ln Z[V,T,\hat{m}]\bigg|_{T=0} = \langle 0|\psi^{+}\sigma_{\mu\nu}\psi|0\rangle_F = \mathrm{Tr}(\tilde{S}\sigma_{\mu\nu}) \tag{49}$$

with

$$\tilde{S} = \frac{1}{\not{P} + i\hat{m} + iM(P)}. \tag{50}$$

We shall consider here only the leading order in the large N_c expansion. Using the Schwinger method [21, 22], we solve the following traces:

$$\begin{aligned} &\mathrm{Tr}\left\{\left(\frac{1}{\not{P} + i(\hat{m} + M(P))} - \frac{1}{\not{P} + i\hat{m}}\right)\sigma_{\mu\nu}\right\} \\ =\ & i\mathrm{Tr}\left\{\left(\frac{(\hat{m} + M(p))}{(p^2 + (\hat{m} + M(p))^2)^2} - \frac{\hat{m}}{(p^2 + \hat{m}^2)^2}\right)\frac{\sigma\cdot F}{2}\sigma_{\mu\nu}\right\} \\ +\ & i\mathrm{Tr}\left\{\frac{1}{(p^2 + (\hat{m} + M(p))^2)^2}\not{p}[\not{P}, M(P)]\sigma_{\mu\nu}\right\}, \end{aligned} \tag{51}$$

where $\sigma\cdot F = \sigma_{\mu\nu}F_{\mu\nu}$. We need to calculate the commutator $[\not{P}, M(P)]$, keeping only terms to order $\mathcal{O}(V)$. It can be easily computed for the dipole-type form factor. Otherwise, we can derive them numerically. Finally, we arrive at the following expression for the magnetic susceptibility:

$$\begin{aligned} \chi_f\langle i\psi_f^\dagger\psi_f\rangle_0 \ =\ & 4N_c\int\frac{d^4p}{(2\pi)^4}\left(\frac{m_f + M_f(p)}{(p^2 + (m_f + M_f(p))^2)^2} - \frac{m_f}{(p^2 + m_f^2)^2}\right) \\ -\ & 4N_c\int\frac{d^4p}{(2\pi)^4}\frac{M_f F(p)F'(p)p}{(p^2 + (m_f + M_f(p))^2)^2}, \end{aligned} \tag{52}$$

where the quark condensate in the chiral limit plays a role of the normalization and is given by:

$$\langle i\psi^\dagger\psi\rangle_0 = 4N_c\int\frac{d^4p}{(2\pi)^4}\frac{M_0F^2(p)}{p^2 + M_0^2F^4(p)} \approx 0.017\,\mathrm{GeV}^3, \tag{53}$$

The infrared region of the first integral in Eq.(52) brings out the contribution of order $\mathcal{O}(m_f \ln m_f)$ which is almost model-independent. In order to calculate the magnetic susceptibility explicitly, we first consider two different regions while integrating: $0 < p < 1\,\mathrm{GeV}$ and $1\,\mathrm{GeV} < p < \infty$. Then we are able to evaluate a part of the free quark analytically. The

---

[1] Our preliminary estimate indicates that the contribution of the meson loops to the dynamic quark mass is not at all small due to the chiral log corrections which are of order $\frac{1}{N_c} m_f \ln m_f$. The meson-loop corrections are even more important than those of order $\mathcal{O}(m_f)$.



FIG. 1: The m_f dependence of the magnetic susceptibility with the form factor of the zero-mode type (Eq. (44)) used. The long-dashed curve draws the contribution of the local vector current, while the short-dashed one depicts that of the nonlocal one. The solid curve represents the total magnetic susceptibility as a function of the current quark mass.

second integral in Eq. (52) is related to the nonlocal contribution without which the vector current is not conserved, i.e. the Ward-Takahashi identity is broken. It arises from the nonlocal quark-quark interaction and is called *nonlocal current*. Summing all contributions, we obtain the final result for the magnetic susceptibility of the QCD vacuum at $q^2 = 0$:

$$\chi_f = \chi_f^{\rm local} + \chi_f^{\rm nonlocal} \tag{54}$$

with

$$\begin{aligned} \chi_f^{\rm local}\langle i\psi_f^\dagger\psi_f\rangle_0 &= \left(29 + 0.103 m_f + \frac{3\,m_f}{2\pi^2}\ln\frac{m_f}{1000\,{\rm MeV}}\right)\ [{\rm MeV}], \qquad (55)\\ \chi_f^{\rm nonlocal}\langle i\psi_f^\dagger\psi_f\rangle_0 &= (18 - 0.095 m_f)\ [{\rm MeV}]. \qquad (56) \end{aligned}$$

In principle, we can also derive the magnetic susceptibility at finite q^2, i.e. the form factor of the magnetic susceptibility of the QCD vacuum.

5. Figure 1 shows the magnetic susceptibility $\chi_f\langle i\psi_f^\dagger\psi_f\rangle_0$ as a function of the current quark mass, with the zero-mode form factor in Eq. (44) employed. It is a monotonically decreasing function as m_f increases, since the third term with $m_f \ln m_f$ in Eq. (55) suppresses the first and second ones noticeably as m_f gets larger. In fact, its value is around -4.0 MeV for up and down quark masses $m_{u,d} \simeq 5$ MeV, whereas for the strange quark mass $m_s \simeq 200$ MeV it turns out to be -49 MeV. Thus, it almost cancels the leading term in the case of the strange magnetic susceptibility χ_s. Moreover, the nonlocal part is responsible for about 40% of the total magnetic susceptibility in the chiral limit and diminishes linearly but slowly as m_f increases. It implies that it is essential to treat the vector current properly in the presence of the nonlocal interaction, in particular, in the case of the magnetic susceptibility. As a result, the magnetic susceptibility for the up and down quarks is: $\chi_{u,d} = 40 \sim 45$ MeV, whereas for the strange quark we obtain $\chi_s = 6 \sim 10$ MeV. As for the up and down quarks, the present result is comparable to that of Ref. [2, 3, 4].



FIG. 2: A comparison between different types of the form factors. The solid curve is drawn with the zero-mode form factor (44), while the long-dashed one depicts the dipole-type form factor. The short-dashed one is for the Gaussian form factor.

A comparison between different types of the form factors is depicted in Fig. 2. As already discussed in Refs. [27, 28, 29], the exponential type of the form factor yields the smallest result as usual. It is interesting to see that the dipole type presents a very similar result for the magnetic susceptibility to that with the zero-mode form factor (44). It is also found that the dependence of the magnetic susceptibility on a type of the form factors is not sensitive.

6. In the present work, we have investigated the magnetic susceptibility of the QCD vacuum within the framework of the nonlocal chiral quark model from the instanton vacuum. We first have constructed the effective partition function in the presence of the external vector and tensor fields as well as the current quark mass. It was shown that for the constant electromagnetic field strength the present result is independent of the path which is selected for the gauge connection. We found that the result is a smoothly decreasing function of m_f: For the up and down quarks $m_{u,d} \simeq 5\,\mathrm{MeV}$ we have $\chi_{u,d}\langle i\psi_{u,d}^\dagger\psi_{u,d}\rangle_0 = 40 \sim 45\,\mathrm{MeV}$ and for the strange quark we obtain $\chi_s\langle i\psi_s^\dagger\psi_s\rangle_0 = 6 \sim 10\,\mathrm{MeV}$. As for the up and down quarks the present results are comparable to the estimate of the vector dominance and QCD sum rule approach [2, 3, 4].

The present work is supported by Korea Research Foundation Grant (KRF-2003-041-C20067). MM acknowledges the support of the Brain Pool program 2004 which makes it possible for him to visit Pusan National University.

[1] V. M. Braun, S. Gottwald, D. Y. Ivanov, A. Schäfer and L. Szymanowski, Phys. Rev. Lett. **89** (2002) 172001 [arXiv:hep-ph/0206305].

[2] V. M. Belyaev and Y. I. Kogan, Yad. Fiz. **40** (1984) 1035.

[3] I. I. Balitsky, A. V. Kolesnichenko and A. V. Yung, Sov. J. Nucl. Phys. **41** (1985) 178 [Yad. Fiz. **41** (1985) 282].

[4] P. Ball, V. M. Braun and N. Kivel, Nucl. Phys. B **649** (2003) 263 [arXiv:hep-ph/0207307].

[5] E. V. Shuryak, Nucl. Phys. B **203** (1982) 93.

[6] D. Diakonov and V. Y. Petrov, Nucl. Phys. B **245** (1984) 259.

[7] M. C. Chu, J. M. Grandy, S. Huang and J. W. Negele, Phys. Rev. D **49** (1994) 6039 [arXiv:hep-lat/9312071].

[8] J. W. Negele, Nucl. Phys. Proc. Suppl. **73** (1999) 92 [arXiv:hep-lat/9810053].

[9] T. DeGrand, Phys. Rev. D **64** (2001) 094508 [arXiv:hep-lat/0106001].

[10] P. Faccioli and T. A. DeGrand, Phys. Rev. Lett. **91** (2003) 182001 [arXiv:hep-ph/0304219].

[11] P. O. Bowman, U. M. Heller, D. B. Leinweber, A. G. Williams and J. b. Zhang, Nucl. Phys. Proc. Suppl. **128** (2004) 23 [arXiv:hep-lat/0403002].

[12] D. Diakonov and V. Y. Petrov, Nucl. Phys. B **272** (1986) 457.

[13] D. Diakonov, M. V. Polyakov and C. Weiss, Nucl. Phys. B **461** (1996) 539 [arXiv:hep-ph/9510232].

[14] D. Diakonov, Prog. Part. Nucl. Phys. **51** (2003) 173 [arXiv:hep-ph/0212026].

[15] C. k. Lee and W. A. Bardeen, Nucl. Phys. B **153** (1979) 210.

[16] T. Schäfer and E. V. Shuryak, Rev. Mod. Phys. **70** (1998) 323 [arXiv:hep-ph/9610451].

[17] M. M. Musakhanov and H.- Ch. Kim, Phys. Lett. B **572** (2003) 181 [arXiv:hep-ph/0206233].

[18] M. M. Musakhanov and F. C. Khanna, Phys. Lett. B **395** (1997) 298 [arXiv:hep-ph/9610418].

[19] E. D. Salvo and M. M. Musakhanov, Eur. Phys. J. C **5** (1998) 501 [arXiv:hep-ph/9706537].

[20] V. Y. Petrov, M. V. Polyakov, R. Ruskov, C. Weiss and K. Goeke, Phys. Rev. D **59** (1999) 114018 [arXiv:hep-ph/9807229].

[21] J. S. Schwinger, Phys. Rev. **82** (1951) 664.

[22] A. I. Vainshtein, V. I. Zakharov, V. A. Novikov and M. A. Shifman, Sov. J. Nucl. Phys. **39** (1984) 77 [Yad. Fiz. **39** (1984) 124].

[23] G. Ripka, *"Quarks bound by Chiral Fields"*, Oxford Science Publications, Oxford (1997) and references therein.

[24] B. Golli, W. Broniowski and G. Ripka, Phys. Lett. B **437** (1998) 24 [arXiv:hep-ph/9807261].

[25] M. Musakhanov, Eur. Phys. J. C **9** (1999) 235 [arXiv:hep-ph/9810295].

[26] M. Musakhanov, Nucl. Phys. A **699** (2002) 340 [arXiv:hep-ph/0104163].

[27] M. Franz, H.-Ch. Kim and K. Goeke, Nucl. Phys. A **699** (2002) 541 [arXiv:hep-ph/9908400].

[28] H. A. Choi and H.-Ch. Kim, Phys. Rev. D **69** (2004) 054004 [arXiv:hep-ph/0308171].

[29] H. J. Lee, C. H. Hyun, C. H. Lee and H.-Ch. Kim, arXiv:hep-ph/0405217.